

## **Libri di testo della Composizione Corporea**

Heyward VH and Stolarczyk LM. Applied Body Composition Assessment. *Human Kinetics*, 1996.

Roche AF, Heymsfield SB, Lohman TG. Human Body Composition. *Human Kinetics*, 1996.

## **Principi sull'Analisi della Composizione Corporea**

Baumgartner RN, et al. Bioelectric impedance for body composition. *Exerc. Spt. Sci. Rev.* 18:193-224, 1990.

Chumlea WC, et al. Bioelectric impedance methods for the estimation of body composition. *Can. J. Spt. Sci.* 15(3):172-179, 1990.

Deurenberg P, et al. Factors affecting bioelectrical impedance measurements in humans. *Eur. J. Clin. Nutr.* 1988; 42:1017-1022.

Ellis J, et. al. Bioelectrical impedance methods in clinical research: a follow-up to the NIH technology assessment conference. *Nutrition*, Volume 15, Nos. 11/12, 1999.

Gallagher DG, et al. Healthy percentage body fat ranges: an approach for developing guidelines based on body mass index. *American Journal of Clinical Nutrition*, 2000; 72:694-701.

Girandola RN, et al. The validity of bioelectrical impedance to predict body composition. Olympic Sci. Congress, *New Horizons of Human Movement*, 1988.

Kyle UG, et al. Fat-Free and Fat Mass Percentiles in 5225 Healthy Subjects Aged 15 to 98 Years. *Nutrition*, 17:534-541, 2001.

Lohman TG. Advances in body composition assessment. *Current Issues in Exercise Science Series, Monograph # 3*, Human Kinetics Publishers, Champaign, IL, 1992.

Lukaski HC, et al. Assessment of fat-free mass using bioelectrical impedance measurements of the human body. *American Journal of Clinical Nutrition*, April 1985; 41:810-817.

Lukaski HC, et al. Validation of tetrapolar bioelectrical impedance methods to assess human body composition. *J. Appl. Physiol.* 60(4):1327-1332, 1986.

Lukaski HC. Methods for the assessment of human body composition: traditional and new. *American Journal of Clinical Nutrition*, 1987; 46:537-56.

Nyboer J. Electrical impedance plethysmography. A physical and physiologic approach to peripheral vascular study. *Circulation*, Vol. 2, Dec. 1950.

Roche AF, et al. Estimation of body composition by impedance. *Med. Sci. Sports Exerc.* 19(2):40, Suppl., Apr. 1987.

Segal KR, et al. Lean body mass estimation by bioelectrical impedance analysis: a four-site cross-validation study. *American Journal of Clinical Nutrition*, 1988; 47:7-14.

Van Loan MD. Bioelectrical impedance analysis to determine fat-free mass, total body water and

body fat. *Sports Med.* 10(4):205-217, 1990.

### **Confronto di Bioimpedenza con altri metodi**

Belford M, et al. The validity of bioelectrical impedance, near-infrared interactance and skinfold equations for estimating body composition in females. *Med. Sci. Sports Exerc.* 25(5):S162, 1993.

Brodie DA, et al. Body fat estimations by electrical impedance and infra-red interactance. *Int. J. Sports Med.* 13(4):319-325, 1992.

Cunningham JJ. New approaches to the noninvasive assessment of body composition: bioelectrical impedance analysis and total body electrical conductivity. *Nutr. Int.* 3:6-10, 1987.

Johnson K, et al. Comparison of circumference, skinfold, bioelectrical impedance and hydrodensitometry to estimate percent body fat during weight loss. *Fed. Proc.* 46(4), Mar. 1987.

Kushner RF, et al. Estimation of lean body mass by bioimpedance analysis compared to skinfold anthropometry. *American Journal of Clinical Nutrition*, 45(4):830, Apr. 1987.

Levenhagen DK, et al. A Comparison of Air Displacement Plethysmography with Three Other Techniques to Determine Body Fat in Healthy Adults. *Journal of Parenteral and Enteral Nutrition*, September-October 1999; Vol. 23, No. 5; 293-299.

Lin KH, et al. Reliability of Body Fat Measurement: Skinfold, Bioelectrical Impedance Analysis and Infrared. *Journal of the Physical Therapy Association of the Republic of China*; September 1995; 102-108.

Meguid MM, et al. Bioelectrical impedance (BIA) method in body composition studies. *J. Parent. Ent. Nutr.* 11(1), Jan-Feb. 1987.

Pasco JA, et al. Body fat estimated from anthropometric and electrical impedance measurement. *Human Nutrition: Clinical Nutrition*, 1985 Sep;39(5):365-9.

Sinning WE, et al. Variability of estimating body composition measures by skinfolds and bioresistance. *Med. Sci. Sports Exerc.* 19(2):39, Supp., Apr. 1987.

Thompson DL, et al. Effects of hydration and dehydration on body composition analysis: a comparative study of bioelectric impedance analysis and hydrodensitometry. *J. Sports Med. Phys. Fitness*, 1991; 31(4):565-70.

Vansant G, et al. Assessment of body composition by skinfold anthropometry and bioelectrical impedance technique: a comparative study. *J. Par. Ent. Nutr.* 1994; 18:427-429.

Villar J, et al. Bioimpedance or anthropometry? *American Journal of Clinical Nutrition*, 1989; 43:129-137.

Wang J, et al. Validation of body fat estimation by BIA using dual photon absorptiometry (DPA). *Fed. Proc.* 1987; 46(4).

## **Obesità**

Bedogni G, et al. Is body mass index a measure of adiposity in elderly women? *Obesity Research*, 2001 Jan;9(1):17-20.

Blackburn GL. Managing obesity in America: An overview. *Advanced Studies in Medicine*, 2002;2(2):40-49.

Fulcher GR, et al. A comparison of measurements of lean body mass derived by bioelectrical impedance, skinfold thickness and total body potassium. A study in obese and non-obese normal subjects. *Scand. J. Clin. Lab. Invest.* 1991; 51(3):245-53.

Guo SS, et al. Body mass index during childhood, adolescence and young adulthood in relation to adult overweight and adiposity: the Fels Longitudinal Study, *International Journal of Obesity and Related Metabolic Disorders*, 2000 Dec;24(12):1628-35.

Heber D, et al. Clinical detection of sarcopenic obesity by bioelectrical impedance analysis. *American Journal of Clinical Nutrition*; 1996; 64 (supplement 3); 472S-477S.

Lauenborg J, et al. The prevalence of the metabolic syndrome in a Danish population of women with previous GDM is 3-fold higher than the general population. *The Journal of Clinical Endocrinology & Metabolism*, 2005, 10.1210/jc. 2004-1713.

Mokdad AH, et al. The spread of the obesity epidemic in the United States, 1991 - 1998. *Journal of American Medical Association*, 1999; 282:1519-1522.

NIH Consensus Conference Statement, Health Implications of Obesity. *Annals of Internal Medicine*, 1985; 103 (6 pt 2):1073-1077.

Regional Office for the Western Pacific of the World Health Organization, the International Association for the Study of Obesity and the International Obesity Task Force. The Asia-Pacific perspective: Redefining obesity and its treatment. *Health Communications Australia Pty Limited*, February 2000.

Rolls BJ, et al. Energy Density But Not Fat Content of Foods Affected Energy Intake in Lean and Obese Women. *American Journal of Clinical Nutrition*; 1999; Vol. 69; 863-871.

## **Sedentarietà**

Blair SN, et al. Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men. *Journal of American Medical Association*, 1995 Apr 12;273(14):1093-8.

Hageman PA, et al. Comparison of two physical activity monitors during a 1-mile walking field test. *Journal of Exercise Physiology Online*, 2004; 7(3):102-110.

Lee CD, Jackson AS, Blair SN. US weight guidelines: Is it also important to consider cardiorespiratory fitness? *Journal of Obesity and Related Metabolic Disorders*, 1998 Aug;22 Supplement 2:S2-7.

## **Soggetti attivi**

Bland JS. The use of complementary medicine for healthy aging. *Alternative Therapies in Health and Medicine*. 1998; 4:42-48.

Evans W, Rosenberg IH. Biomarkers. *Fireside*, 1992.

Guo SS, et al. Aging, body composition, and lifestyle: the Fels Longitudinal Study. *American Journal of Clinical Nutrition*, 1999; 70: 405-11.

Kyle UG, et al. Physical activity and fat-free and fat mass by bioelectrical impedance in 3853 adults. *Medicine and Science in Sports and Exercise*, 2001;33:576-584.

## **Atleti**

Chapman R, et al. Body composition testing of athletes in the field using bioelectric impedance analysis. *Alaska Med.*, 1992 Apr-Jun;34(2):87-90,95.

Fornetti WC, et al. Reliability and validity of body composition measures in female athletes. *Journal of Applied Physiology*, 1999 Sep;87(3):1114-22.

Johnson GO, et al. The validity of bioelectrical impedance, near-infrared interactance and skinfold equations for estimating percent fat in female gymnast. *Med. Sci. Sports Exerc.* 25(5):S162, 1993.

Lukaski HC, et al. Body composition assessment of athletes using bioelectrical impedance measurements. *J. Sports Med. Phys. Fitness* 1990; 30:434-40.

Mattar R, et al. Body mass index (BMI) as an indicator of body composition: evidence through bioelectrical impedance. *XVIII Simp. Int. Ciênc. Esp.*, Oct. 1992.

Oppliger RA, et al. Bioelectrical impedance prediction of fat-free mass for high school wrestlers validated. *Medicine and Science in Sports and Exercise*, 1991, 23, S73 (Abstract).

Position of the American Dietetic Association, Dietitians of Canada, and the American College of Sports Medicine. Nutrition and athletic performance. *Journal of American Dietetic Association*, 2000; 100:1543-1556.

## **Anziani**

Baumgartner RN, et al. Body Composition in elderly people: Effect of criterion estimates on predictive equations. *American Journal of Clinical Nutrition*, 1991; 53, 1-9.

Berger VA, et al. Reproducibility of body composition and body water spaces measurements in healthy elderly individuals. *The Journal of Nutrition, Health & Aging*, 2000;4(4):24305.

Boeckner LS, et al. Eating behaviors and health history of rural midlife to older women in the Midwestern United States. *Journal of the American Dietetic Association*, 2007; 107: 306-310.

Hughes VA, Evans WJ. Assessment of fat-free mass in an older population using bioelectrical impedance. *Federation Proceedings*, 46(4), 1987.

Omran ML, Morley JE. Assessment of protein energy malnutrition in older persons, part I: History, examination, body composition, and screening tools. *Nutrition*, 2000 Jan;16(1):50-63.

Roubenoff R, et al. Nutrition assessment in long-term care facilities. *Nutrition Review*, 1996 Jan;54(1 Pt 2):S40-2.

Svendsen OL, et al. Measurement of body fat in elderly subjects by dual-energy x-ray absorptiometry, bioelectrical impedance and anthropometry. *American Journal of Clinical Nutrition*, 1991; 53:1117-23.

## **Bambini**

Goran MI, et al. Estimating body composition of young children by using bioelectrical resistance. *J. Appl. Physiol.* 75(4):1776-1780, 1993.

Houtkooper LB, et al. Validity of whole body bioelectrical impedance analysis for body composition assessment in children. *Med. Sci. Sports Exerc.* 19(2):39, Suppl., Apr. 1987.

Houtkooper LB, et al. Validity of bioelectric impedance for body composition assessment in children. *J. Appl. Physiol.* 66(2):814-21, 1989.

Houtkooper LB, et al. Bioelectrical impedance estimation of fat-free mass in children and youth: a cross-validation study. *J. Appl. Physiol.* 72(1):366-373, 1992.

Mazariegos M, et al. Bioelectrical impedance used in association with anthropometry: experience in a field survey of underprivileged Guatemalan children. *Age & Nutr.* 5(2), 1994.

Roche AF, et al. Fat-free mass in children and young adults predicted from bioelectric impedance and anthropometric variables. *American Journal of Clinical Nutrition*, 1989; 50:435-43.

## **Supplementi**

Antonio JA, et al. Effects of a Standardized Guggulsterone Phosphate Supplement on Body Composition in Overweight Adults: A Pilot Study. *Current Therapeutic Research*; April 1999; Vol. 60, No. 4; 220-227.

Colker CM, et al. Effects of Citrus aurantium Extract, Caffeine, and St. John's Wort on Body Fat Loss, Lipid Levels, and Mood States in Overweight Healthy Adults. *Current Therapeutic Research*; March 1999; Vol. 60, No. 3; 145-153.

Kalman D, et al. Effects of Pyruvate Supplementation on Body Composition and Mood. *Current Therapeutic Research*; November 1998; Vol. 59, No. 11; 793-802.

Kalman D, et al. The Effects of Pyruvate Supplementation on Body Composition in Overweight Individuals. *Nutrition*; May 1999; Vol. 15, No. 5; 337-340.

Kalman D, et al. Randomized Trial of an Isocaloric High Protein Diet vs. an Isocaloric Moderate Protein Diet for Short-Term Weight Loss in Healthy Overweight Adults. *The Bariatrician*; Winter 2000/2001; 10-13.

Udani J, et al. Blocking carbohydrate absorption and weight loss: a clinical trial using phase 2 brand proprietary fractionated white bean extract. *Alternative Medicine Review*, 2004; 9(1):63-69.

### **Analisi dei liquidi**

Chertow GM, et al. Bioelectrical impedance analysis predicts survival in hemodialysis patients, abstract #T164, A0969, page 1442. *Journal of the American Society of Nephrology*, September 1996, Volume 7, #9.

Chertow GM, et al. Nutritional assessment with bioelectrical impedance analysis in maintenance hemodialysis patients. *Journal of the American Society of Nephrology*, 1995, 6:75-81.

Cooper BA, et al. Comparing different methods of assessing body composition in end-stage renal failure. *Kidney International*, 2000 Jul;58(1):408-16.

de Vries PMJM, et al. Measurement of transcellular fluid shift during hemodialysis. *Med. & Biol. Eng. & Comput.* 1989; 27:152-158.

Espejo MGA, et al. Determination of extracellular fluid volume using impedance measurements. *Crit. Care Med.* 1989; 17(4):360-363.

Johnson HL, et al. Use of bioelectrical measurements for the estimation of total body water and extracellular space in man. *Fed. Proc.* 46(4), Mar. 1987.

Kushner RF, Schoeller DA. Estimation of total body water by bioelectrical impedance analysis. *American Journal of Clinical Nutrition*, 44: Sept. 1986, 417-424.

Leiter LA, et al. Use of bioelectrical impedance analysis measurement in patients with diabetes. *American Journal of Clinical Nutrition*, 1996; 64: 515S-518S.

Ljungqvist O, et al. Whole body impedance measurements reflect total body water changes. A study in hemodialysis patients. *Swe. Int. J. Clin. Monit. Comput.* 1990; 7(3):163-169.

Lukaski HC, et al. Estimation of body fluid volumes using tetrapolar bioelectrical impedance measurements. *Aviat. Space Environ. Med.* 1988; 59:1163-9.

Lukaski HC. Validation of body composition assessment techniques in the dialysis population. *ASAIO J* 1997, May-June; 43(3):251-5.

Maehara T, et al. Perioperative monitoring of total body water by bioelectrical impedance in children undergoing open heart surgery. *Eur. J. Cardiothorac. Surg.* 1991; 5(5):258-65.

Mayfield SR, et al. Measurement of extracellular water (ECW) in low birth weight infants using bioelectrical reactance. *Am. Ped. Soc. & Soc. Ped. Res.* 1987.

McCullough AJ, et al. Measurements of total body and extracellular water in cirrhotic patients with and without ascites. *Hepatology*, 1991; 14(6):1102-11.

Meguid MM, et al. Rapid bedside method to assess changes in postoperative fluid status with bioelectrical impedance analysis. *Surgery* 1992; 112:502-8.

Scanforla F, et al. On-line bioelectric impedance during hemodialysis: monitoring of body fluids and cell membrane status. *Nephrol. Dial. Transplant. Suppl.* 1 (1990):167-170.

Segal KR, et al. Estimation of body water distribution by bioelectrical impedance. *Fed. Proc.* 1987; 46:1334.

Stall SH. Comparison of five body-composition methods in peritoneal dialysis patients. *American Journal of Clinical Nutrition*, 1996; August; 64(2):125-30.

Van Loan MD, et al. Use of bioimpedance spectroscopy (BIS) to determine extracellular fluid (ECF), intracellular fluid (ICF), total body water (TBW) and fat-free mass (FFM). *Int. Symp. In Vivo Body Comp. Studies*, Texas, Nov. 1992.

Westendorp RGJ, et al. Weight changes in critically ill patients evaluated by fluid balances and impedance measurements. *Crit. Care Med.* 1993; 21(6):871-877.

Wu TJ, et al. Bioelectrical impedance analysis of nutritional status in uremic patients on regular hemodialysis. *Taiwan I. Hsueh. Hui. Tsa. Chin.* 1991; 90(11):1044-8.

### **Massa cellulare corporea**

De Lorenzo A, et al. Predicting body cell mass with bioimpedance by using theoretical methods: a technological review. *Journal of Applied Physiology*, 82(5): 1542-1558, 1997.

Jacobs DO. Bioelectrical impedance analysis: a way to assess changes in body cell mass in patients with acquired immunodeficiency syndrome. *J. Parent. Ent. Nutr.*, 17(5):401-2, 1993.

Kotler DP, et al. Prediction of body cell mass, fat-free mass, and total body water with bioelectrical impedance analysis: effects of race, sex, and disease, *American Journal of Clinical Nutrition*, 1996; 64 (supplement); 489S-497S.

Schloerb PR, et al. Bioimpedance as a measure of total body water and body cell mass in surgical nutrition. *Eur. Surg. Res.* 18(SL):3, 1986.